

NSIPS: Nonlinear Semi-Infinite Programming Solver

A. Ismael F. Vaz

Edite M.G.P. Fernandes

Production and Systems Department
Engineering School

Minho University

Braga - Portugal

{aivaz, emgpf}@dps.uminho.pt

M. Paula S.F. Gomes

Mechanical Engineering Department
Imperial College of Science,
Technology and Medicine

London SW7 2BX - UK

p.gomes@ic.ac.uk

18-20 June, 2004

Talk partially financed by the Portuguese Calouste Gulbenkian Foundation

Contents

- Semi-Infinite Programming (SIP)
- Nonlinear SIP Solver - (NSIPS)
 - ★ Discretization method
 - ★ Sequential quadratic programming method
 - ★ Constraint transcription methods
 - * Interior point method
 - * Penalty (multiplier) method
- Numerical results / Conclusions

Semi-Infinite Programming

$$\begin{aligned}
 & \min_{x \in R^n} f(x) \\
 \text{s.t. } & g_i(x, t) \leq 0, \quad i = 1, \dots, m \\
 & h_i(x) \leq 0, \quad i = 1, \dots, o \\
 & h_i(x) = 0, \quad i = o + 1, \dots, q \\
 & \forall t \in T
 \end{aligned} \tag{1}$$

where $f(x)$ is the objective function, $h_i(x)$ are the finite constraint functions, $g_i(x, t)$ are the infinite constraint functions and $T \subset R^p$ is, usually, a cartesian product of intervals $([\alpha_1, \beta_1] \times [\alpha_2, \beta_2] \times \cdots \times [\alpha_p, \beta_p])$.

Solver

NSIPS (Nonlinear Semi-Infinite Programming Solver)

Solver

NSIPS (Nonlinear Semi-Infinite Programming Solver)

Version 2.1 publicly available in the internet

<http://www.norg.uminho.pt/aivaz/>, implementing the four methods in a total of seven algorithms.

Solver

NSIPS (Nonlinear Semi-Infinite Programming Solver)

Version 2.1 publicly available in the internet

<http://www.norg.uminho.pt/aivaz/>, implementing the four methods in a total of seven algorithms.

NSIPS receives the problem in the (SIP)AMPL format.

Solver

NSIPS (Nonlinear Semi-Infinite Programming Solver)

Version 2.1 publicly available in the internet

<http://www.norg.uminho.pt/aivaz/>, implementing the four methods in a total of seven algorithms.

NSIPS receives the problem in the (SIP)AMPL format.

The method is select by the *nsips_options*, and there are many other options for each method.

Solver

NSIPS (Nonlinear Semi-Infinite Programming Solver)

Version 2.1 publicly available in the internet

<http://www.norg.uminho.pt/aivaz/>, implementing the four methods in a total of seven algorithms.

NSIPS receives the problem in the (SIP)AMPL format.

The method is select by the *nsips_options*, and there are many other options for each method.

The NPSOL software is used to solve the finite subproblems (discretization and SQP methods).

Solver

NSIPS (Nonlinear Semi-Infinite Programming Solver)

Version 2.1 publicly available in the internet

<http://www.norg.uminho.pt/aivaz/>, implementing the four methods in a total of seven algorithms.

NSIPS receives the problem in the (SIP)AMPL format.

The method is select by the *nsips_options*, and there are many other options for each method.

The NPSOL software is used to solve the finite subproblems (discretization and SQP methods).

NSIPS is available in the NEOS server (www-neos.mcs.anl.gov).

Discretization method - Three versions

A sequence of finite problems are solved. The finite problems are obtained from the SIP problem where the infinite constraints are evaluated at a finite set of points $\tilde{T}[h^k] \subseteq T[h^k]$, where $T[h^k] \subseteq T$ is a uniform grid of points with space h^k .

Versions adapted for nonlinear SIP and implemented:

- Hettich (1986, 1990)
- Reemtsen (1991)
- Hettich with pseudo-number generation.

Discretization method

- STEP 0: Define $T[h^0]$. Let $\tilde{T}[h^0] = T[h^0]$. Solve the $\text{NLP}(\tilde{T}[h^0])$ and let x_0 be the solution found.
- STEP k : If x_{k-1} is not feasible for all the points in the set $T[h^{k-1}]$
 - ★ THEN: Insert all the infeasible points in the set $\tilde{T}[h^{k-1}]$. Solve the $\text{NLP}(\tilde{T}[h^{k-1}])$ and let x_{k-1} be the solution found. Continue with step k .
 - ★ ELSE: If the maximum number of refinements is reached then stop. Else build the set $\tilde{T}[h^k]$ from $T[h^k]$ and $\tilde{T}[h^{k-1}]$. Solve the $\text{NLP}(\tilde{T}[h^k])$ and let x_k be the solution found. Go to step $k + 1$.

Reduced problem

Problem with no finite constraints and only one infinite variable.

$$\begin{aligned} & \min_{x \in R^n} f(x) \\ s.t. \quad & g_i(x, t) \leq 0, \quad i = 1, \dots, m \\ & \forall t \in T \equiv [a, b] \end{aligned} \tag{2}$$

Sequential Quadratic Programming

Considering the reduced problem (2), the sequential quadratic programming is based on the quadratic semi-infinite programming (QSI)

$$\begin{aligned} \min_{d \in R^n} f_Q(d) &= \frac{1}{2} d^T H_k d + d^T \nabla f(x_k) \\ s.t. \quad d^T \nabla_x g_i(x_k, t) + g_i(x_k, t) &\leq 0, \\ i &= 1, \dots, m, \quad \forall t \in [a, b] \end{aligned}$$

where H_k is an approximation to $\nabla_{xx}^2 \mathcal{L}(x_k, v)$.

SQP

The solution of the QSI problem is d_k and

$$x_{k+1} = x_k + \alpha_k d_k, \quad k = 1, 2, \dots$$

$\{x_k\} \rightarrow x^*$, solution to the initial SIP problem.

The Lagrangian of the QSI problem is

$$\begin{aligned} \mathcal{L}_Q(d, v) &= \frac{1}{2} d^T H_k d + d^T \nabla f(x_k) \\ &\quad + \sum_{j=1}^m \int_a^b \left(d^T \nabla_x g_j(x_k, t) + g_j(x_k, t) \right) dv_j(t) \end{aligned}$$

Solving the QSI


The dual problem $\min_{v \in \mathcal{V}^*} \mathcal{L}_Q^*(v) \equiv -\mathcal{L}_Q(d(v), v)$ is solved by a linear parametrization of the dual variables.

$$v_j(t) = \begin{cases} w_{1j}(t - a), & \text{for } t \in [a, t_1); \\ a_{ij} + w_{i+1j}(t - t_i), & \text{for } t \in [t_i, t_{i+1}), \\ & i = 1, 2, \dots, l-1; \\ a_{lj} + w_{l+1j}(t - t_l), & \text{for } t \in [t_l, b]; \end{cases}$$

$j = 1, \dots, m$, where

$$a_{ij} = \sum_{p=1}^i h_{pj} + \sum_{p=1}^i w_{pj}(t_p - t_{p-1}), \quad i = 1, \dots, l$$

Example with $m = 2, l = 2$



w are the linear segments slope, h are the jumps and t are the discontinuity points.

Merit function

$$\phi(x, \mu) = f(x) + \frac{1}{2}\mu \sum_{i=1}^m \int_a^b [g_i(x, t)]_+^2 dt$$

where $[z]_+ = \max\{0, z\}$.

A strategy for computing the penalty parameter.

Numerical integrals computation - Numerical adaptative formulae (Gaussian or trapezoid).

SQP - Dual method

1. Given x_0 . Let $k = 0$ and $H_0 = I$.
2. Compute H_k using a BFGS quasi-Newton updating formula.
3. Solve the QSI problem to obtain the search direction d_k .
4. If $d_k = \mathbf{0}$ then stop.
5. Find α_k such that $x_{k+1} = x_k + \alpha_k d_k$ sufficiently decreases the merit function.
6. If there is not a major difference between x_{k+1} and x_k then stop with x_{k+1} as an approximated solution. Otherwise do $k = k + 1$ and go to step 2.

Constraint transcription

Considering the reduced problem (2), the infinite constraints $g_i(x, t) \leq 0, \forall t \in T$, are transformed into $\int_T [g_i(x, t)]_+ dt = 0$ where $[z]_+ = \max\{0, z\}$.

The SIP is then transformed into

$$\begin{aligned} & \min_{x \in R^n} f(x) \\ s.t. \quad & G_i(x) \equiv \int_T [g_i(x, t)]_+ dt = 0 \\ & i = 1, \dots, m \end{aligned}$$

Constraint functions not differentiable.

Approximate problem

$$\min_{x \in R^n} f(x)$$

$$s.t. \quad G_{i,\epsilon}(x) \equiv \int_T g_{i,\epsilon}(x, t) dt = 0$$

$$i = 1, \dots, m$$

with $\epsilon \rightarrow 0$ and

$$g_{i,\epsilon}(x, t) = \begin{cases} 0, & \text{if } g_i(x, t) < -\epsilon; \\ \frac{(g_i(x, t) + \epsilon)^2}{4\epsilon}, & \text{if } -\epsilon \leq g_i(x, t) \leq \epsilon; \\ g_i(x, t), & \text{if } g_i(x, t) > \epsilon, \end{cases}$$

Once differentiable constraint functions.

Penalty method

A sequence of subproblems is solved, parameterized by μ

$$\min_{x \in R^n} \phi_S(x, \mu)$$

for a sequence of increasing $\mu > 0$ values.

Simple penalty functions

$$\phi_S^1(x, \mu) = f(x) + \mu \sum_{i=1}^m \int_T g_{i,\epsilon}(x, t) dt$$

Simple penalty functions

$$\phi_S^1(x, \mu) = f(x) + \mu \sum_{i=1}^m \int_T g_{i,\epsilon}(x, t) dt$$

$$\phi_S^2(x, \mu) = f(x) + \frac{\mu}{2} \sum_{i=1}^m \int_T g_{i,\epsilon}(x, t)^2 dt$$

Simple penalty functions

$$\phi_S^1(x, \mu) = f(x) + \mu \sum_{i=1}^m \int_T g_{i,\epsilon}(x, t) dt$$

$$\phi_S^2(x, \mu) = f(x) + \frac{\mu}{2} \sum_{i=1}^m \int_T g_{i,\epsilon}(x, t)^2 dt$$

and

$$\phi_S^3(x, \mu) = f(x) + \mu \sum_{i=1}^m \int_T \left(e^{g_{i,\epsilon}(x, t)} - 1 \right) dt$$

Relaxed problem to satisfy LICQ

$$\min_{x \in R^n} f(x)$$

$$s.t. \quad G_{i,\epsilon}(x) \leq \tau$$

$$i = 1, \dots, m$$

$$\tau > 0 \quad (\tau(\epsilon) \rightarrow 0)$$

Multiplier method

A sequence of subproblems is solved

$$\min_{x \in R^n} \phi_{AL}(x, \lambda, \mu)$$

where ϕ_{AL} is the augmented Lagrangian penalty function

Multiplier method

A sequence of subproblems is solved

$$\min_{x \in R^n} \phi_{AL}(x, \lambda, \mu)$$

where ϕ_{AL} is the augmented Lagrangian penalty function

$$\begin{aligned} \phi_{AL}(x, \lambda, \mu) = & f(x) + \sum_{i=1}^m \lambda_i \left(\int_T g_{i,\epsilon}(x, t) dt - \tau \right) \\ & + \frac{\mu}{2} \sum_{i=1}^m \left(\int_T g_{i,\epsilon}(x, t) dt \right)^2 \end{aligned}$$

where $\lambda = (\lambda_1, \dots, \lambda_m)^T$ is the Lagrange multipliers vector.

Lagrange multipliers update

Since the optimum Lagrange multipliers are unknown before computing the solution, an updating formula for the Lagrange multipliers is used.

$$\lambda_i^{k+1} = \lambda_i^k + \mu^k \int_T g_{i,\epsilon}(x^k, t) dt, \quad i = 1, \dots, m.$$

Multiplier method

A sequence of subproblems is solved

$$\min_{x \in R^n} \phi_E(x, \lambda, \mu)$$

where ϕ_E is the exponential penalty function

Multiplier method

A sequence of subproblems is solved

$$\min_{x \in R^n} \phi_E(x, \lambda, \mu)$$

where ϕ_E is the exponential penalty function

$$\phi_E(x, \lambda, \mu) = f(x)$$

$$+ \frac{1}{\mu} \sum_{i=1}^m \lambda_i \left(e^{\mu(\int_T g_{i,\epsilon}(x,t) dt - \tau)} - 1 \right)$$

Multiplier method

A sequence of subproblems is solved

$$\min_{x \in R^n} \phi_E(x, \lambda, \mu)$$

where ϕ_E is the exponential penalty function

$$\phi_E(x, \lambda, \mu) = f(x)$$

$$+ \frac{1}{\mu} \sum_{i=1}^m \lambda_i \left(e^{\mu(\int_T g_{i,\epsilon}(x,t) dt - \tau)} - 1 \right)$$

An updating formula for the Lagrange multipliers is used

$$\lambda_i^{k+1} = \lambda_i^k e^{\mu^k (\int_T g_{i,\epsilon}(x^k, t) dt - \tau)}, \quad i = 1, \dots, m$$

Multiplier penalty framework

1. Given an initial guess for x and λ , and parameters μ , ϵ and $\tau(\epsilon)$.
2. Exterior iteration. The initial guess for the interior iterations is the last approximation computed.
3. Interior iterations. For μ and λ , solve the unconstrained problem

$$\min_{x \in R^n} \phi(x, \lambda, \mu)$$

through a BFGS quasi-Newton technique and a line search with an Armijo like rule that significantly reduces the penalty function.

Solution: $x^*(\mu)$.

4. If the computed approximation is infeasible ($\int_T g_{i,\epsilon}(x^*(\mu), t) dt - \tau > 0$, $i = 1, \dots, m$) then update the penalty parameter μ , the multipliers vector λ and proceed with another exterior iteration.
5. Otherwise, if there is a significant evolution from the last two approximations computed for different differentiable parameters (ϵ e $\tau(\epsilon)$) then update the differentiability parameter and proceed with another exterior iteration.
6. Stop with the last computed approximation being an approximation to the SIP solution ($x^* \leftarrow x^*(\mu)$).

Primal-dual interior point method

From the relaxed problem, the barrier problem is formed by placing the slack variables in the barrier term

$$\begin{aligned} \min_{x \in R^n, s \in R^m} \quad & f(x) - \mu \sum_{i=1}^m \log(s_i + \tau) \\ \text{s.t.} \quad & \int_T g_{i,\epsilon}(x, t) dt + s_i = 0, \quad i = 1, \dots, m \end{aligned}$$

with $g_{i,\epsilon}(x, t) = \frac{g_i(x, t) + \sqrt{g_i(x, t)^2 + \epsilon^2}}{2}$ and $\epsilon \rightarrow 0$ ($\epsilon > 0$).

The barrier problem is solved for a sequence of $\mu(\rightarrow 0)$ values.

By applying the Newton method to the first order KKT system:

Newton system

$$\begin{pmatrix} H & 0 & J \\ 0 & \Lambda & S \\ -J^T & -I & 0 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta s \\ \Delta \lambda \end{pmatrix} = \begin{pmatrix} \sigma \\ \gamma \\ \rho \end{pmatrix}$$

with

$$H = \nabla^2 f - \sum_{i=1}^m \lambda_i \int_T \nabla_{xx}^2 g_{i,\epsilon}(x, t) dt$$

Newton system

$$\begin{pmatrix} H & 0 & J \\ 0 & \Lambda & S \\ -J^T & -I & 0 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta s \\ \Delta \lambda \end{pmatrix} = \begin{pmatrix} \sigma \\ \gamma \\ \rho \end{pmatrix}$$

with

$$H = \nabla^2 f - \sum_{i=1}^m \lambda_i \int_T \nabla_{xx}^2 g_{i,\epsilon}(x, t) dt$$

$$J = \left(\int_T \nabla_x g_{1,\epsilon}(x, t) dt, \dots, \int_T \nabla_x g_{m,\epsilon}(x, t) dt \right)$$

Newton system

$$\begin{pmatrix} H & 0 & J \\ 0 & \Lambda & S \\ -J^T & -I & 0 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta s \\ \Delta \lambda \end{pmatrix} = \begin{pmatrix} \sigma \\ \gamma \\ \rho \end{pmatrix}$$

with

$$H = \nabla^2 f - \sum_{i=1}^m \lambda_i \int_T \nabla_{xx}^2 g_{i,\epsilon}(x, t) dt$$

$$J = \left(\int_T \nabla_x g_{1,\epsilon}(x, t) dt, \dots, \int_T \nabla_x g_{m,\epsilon}(x, t) dt \right)$$

$$S = \text{diag}(s_i + \tau), \quad \Lambda = \text{diag}(\lambda_i), \quad i = 1, \dots, m$$

Newton system

$$\begin{pmatrix} H & 0 & J \\ 0 & \Lambda & S \\ -J^T & -I & 0 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta s \\ \Delta \lambda \end{pmatrix} = \begin{pmatrix} \sigma \\ \gamma \\ \rho \end{pmatrix}$$

with

I = Identity matrix

Newton system

$$\begin{pmatrix} H & 0 & J \\ 0 & \Lambda & S \\ -J^T & -I & 0 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta s \\ \Delta \lambda \end{pmatrix} = \begin{pmatrix} \sigma \\ \gamma \\ \rho \end{pmatrix}$$

with

I = Identity matrix

$\sigma = -\nabla f - J\lambda$ (Dual infeasibility)

Newton system

$$\begin{pmatrix} H & 0 & J \\ 0 & \Lambda & S \\ -J^T & -I & 0 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta s \\ \Delta \lambda \end{pmatrix} = \begin{pmatrix} \sigma \\ \gamma \\ \rho \end{pmatrix}$$

with

I = Identity matrix

$\sigma = -\nabla f - J\lambda$ (Dual infeasibility)

$\gamma = \mu e - S\Lambda e$

Newton system

$$\begin{pmatrix} H & 0 & J \\ 0 & \Lambda & S \\ -J^T & -I & 0 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta s \\ \Delta \lambda \end{pmatrix} = \begin{pmatrix} \sigma \\ \gamma \\ \rho \end{pmatrix}$$

with

I = Identity matrix

$\sigma = -\nabla f - J\lambda$ (Dual infeasibility)

$\gamma = \mu e - S\Lambda e$

$\rho = \bar{g} + s$ (Primal infeasibility)

Newton system

$$\begin{pmatrix} H & 0 & J \\ 0 & \Lambda & S \\ -J^T & -I & 0 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta s \\ \Delta \lambda \end{pmatrix} = \begin{pmatrix} \sigma \\ \gamma \\ \rho \end{pmatrix}$$

with

I = Identity matrix

$\sigma = -\nabla f - J\lambda$ (Dual infeasibility)

$\gamma = \mu e - S\Lambda e$

$\rho = \bar{g} + s$ (Primal infeasibility)

$\bar{g} = (G_{1,\epsilon}, \dots, G_{m,\epsilon})^T$

Newton system

$$\begin{pmatrix} H & 0 & J \\ 0 & \Lambda & S \\ -J^T & -I & 0 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta s \\ \Delta \lambda \end{pmatrix} = \begin{pmatrix} \sigma \\ \gamma \\ \rho \end{pmatrix}$$

$(\Delta x, \Delta s, \Delta \lambda)$ is the Newton direction and

$$x_{k+1} = x_k + \alpha_k \Delta x_k$$

$$s_{k+1} = s_k + \alpha_k \Delta s_k$$

$$\lambda_{k+1} = \lambda_k + \alpha_k \Delta \lambda_k$$

Implemented merit functions

Choosing α to obtain feasibility and convergence to the minimum.

$$\phi(x, s; \mu, \beta) = f(x) - \mu \sum_{i=1}^m \log(s_i + \tau) + \frac{\beta}{2} \rho^T \rho$$

Implemented merit functions

Choosing α to obtain feasibility and convergence to the minimum.

$$\phi(x, s; \mu, \beta) = f(x) - \mu \sum_{i=1}^m \log(s_i + \tau) + \frac{\beta}{2} \rho^T \rho$$

$$\begin{aligned} \mathcal{L}_A(x, s, \lambda; \mu, \beta) = & f(x) - \mu \sum_{i=1}^m \log(s_i + \tau) + \lambda^T \rho \\ & + \frac{\beta}{2} \rho^T \rho \end{aligned}$$


Quasi-Newton strategy with an approximation to the Hessian of the Lagrangian.

Primal-dual interior point algorithm

1. Given x_0 , ϵ , τ , θ , δ_μ and δ_f .
2. Compute $s_{i,0}$ and $\lambda_{i,0}$, $i = 1, \dots, m$. Let $k = 0$.
3. Let $y_{eps} = x_k$ the last y computed for a given ϵ .
4. Compute or update μ_k .
5. Stopping criteria. If the stopping criteria is verified then if there is a significant difference between y_{eps} and x_k reduce ϵ , τ , update the slack variables and go to step 3; Otherwise stop.
6. Update B_k by a BFGS formula. If $k = 0$ then B_k = Identity matrix.

7. Solve the KKT system to obtain the search direction $(\Delta x_k, \Delta s_k, \Delta \lambda_k)$.
8. Compute β and α_{max} .
9. Compute α_k , using a strategy that significantly reduce the merit function
10. Compute x_{k+1} , s_{k+1} and λ_{k+1} .
11. Go to step 4.

Numeric integration



Numerical results / Conclusions

- *Discretization method*

- ★ Solves all problems in the (SIP)AMPL database (over 160 problems) except problems *elke2* and *blankenship2/3*;
- ★ Solution found in the finest grid (no KKT point);
- ★ Needs NPSOL to solve the finite subproblems.

Numerical results / Conclusions

- *Discretization method*
 - ★ Solves all problems in the (SIP)AMPL database (over 160 problems) except problems *elke2* and *blankenship2/3*;
 - ★ Solution found in the finest grid (no KKT point);
 - ★ Needs NPSOL to solve the finite subproblems.
- *SQP method*
 - ★ Solves all problems with only one infinite variable and without finite constraints, except for the robotics problems;
 - ★ Needs NPSOL to solve the finite subproblems.

Numerical results (cont.) / Conclusions

- *Penalty method*
 - ★ Solves all problems with only one infinite variable and without finite constraints;

Numerical results (cont.) / Conclusions

- *Penalty method*
 - ★ Solves all problems with only one infinite variable and without finite constraints;
- *Interior point method*
 - ★ Solves 75% of problems with only one infinite variable and without finite constraints.

The End

email: aivaz@dps.uminho.pt

emgpf@dps.uminho.pt

p.gomes@ic.ac.uk

Web <http://www.norg.uminho.pt/aivaz/>

First Page